



**The Trusted  
Choice.....**

- In Prevention of Stroke in Patients with NVAF  
&
- In Treatment of VTE and PE

Rx

# Apiquis™

2.5  
5

Apixaban 2.5/5mg Tablets



**Rx**

# Ap quis <sup>TM</sup>

**2.5**  
**5**

Apixaban 2.5/5mg Tablets

## Product Description

### Ap quis (Apixaban) Tablets

Available in 2.5 mg & 5 mg strengths

#### Indications:

- ◆ Stroke and Systemic Embolism Prevention in NVAF

Ap quis is used to reduce the risk of stroke and systemic embolism in patients with nonvalvular atrial fibrillation (NVAF), a condition that increases the likelihood of blood clots forming and traveling to vital organs like the brain.

- ◆ Prevention of Deep Vein Thrombosis (DVT)

It is also indicated for the prevention of deep vein thrombosis (DVT), a condition where blood clots form in the deep veins, particularly in the legs, which can then travel to the lungs and cause a pulmonary embolism (PE).

- ◆ Treatment of DVT and PE & Recurrence Prevention

Ap quis is used in the treatment of DVT and PE, helping to dissolve the existing clots. In addition, it reduces the risk of future clot formation following the initial treatment phase.

#### Mechanism of Action:

- ◆ Ap quis is a oral anticoagulant (NOAC) and works by selectively inhibiting Factor Xa, an enzyme crucial in the blood clotting process. It functions by:
  - ◆ Inhibiting both free and clot-bound Factor Xa, as well as prothrombinase activity, which prevents the formation of thrombin (an enzyme that promotes clotting).
  - ◆ No effect on platelet aggregation: Unlike some other anticoagulants, Ap quis does not directly affect the aggregation of platelets, which are involved in the formation of blood clots.
  - ◆ Thrombus formation prevention: By inhibiting Factor Xa, Ap quis decreases the generation of thrombin, which ultimately reduces the formation of blood clots and the risk of thromboembolic events (such as stroke, DVT, and PE).

#### Recommended Dosage:

- ◆ For DVT Prevention : 2.5 mg taken twice a day or as prescribed by the doctor
- ◆ For Nonvalvular Atrial Fibrillation (NVAF): 5 mg taken twice a day or as prescribed by the doctor
- ◆ For DVT & PE Treatment : 5 mg taken twice a day or as prescribed by the doctor



Corp. & Regd. Office:

1009-1010, Chiranjiv Tower, 43, Nehru Place,  
New Delhi - 110019 Tel.: 011-47589500-51(50 lines)  
Web: [www.medicamen.com/lifesciences](http://www.medicamen.com/lifesciences)  
E-mail: [info@medicamenlifesciences.com](mailto:info@medicamenlifesciences.com)

I am: \_\_\_\_\_  
 Call me on: \_\_\_\_\_  
 Mail me at: \_\_\_\_\_

2025 All rights reserved, Medicamen Lifesciences Pvt Ltd